

LAMB & KID MILK REPLACER WITH CITRISTIM MEDICATED

For the prevention of coccidiosis in **young sheep** caused by *Eimeria bakuensis* and *Eimeria crandallis*, *Eimeria ovinoidalis* and *Eimeria parva*. For the prevention of coccidiosis in **young goats** caused by *Eimeria christensenii* and *Eimeria ninakohlyakimovae*. Feed at least 28 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

ACTIVE DRUG INGREDIENT

Decoquinatate..... 18.16 g/ton

GUARANTEED ANALYSIS

Crude Protein, min..... 24.00%
 Crude Fat, min 30.00%
 Crude Fiber, max..... 0.15%
 Ash, max..... 6.50%
 Calcium, min..... 0.95%
 Calcium, max..... 1.45%
 Phosphorus, min 0.80%
 Selenium (added), max 0.3 ppm
 Vitamin A, min..... 30,000 IU/lb
 Vitamin D₃, min..... 5,500 IU/lb
 Vitamin E, min 150 IU/lb

INGREDIENTS

Dried Skimmed Milk, Dried Whey Protein Concentrate, Sodium Propionate, Citric Acid, Calcium Propionate (Preservatives), Dried Whey Product, Dried Milk Protein, Dried Whey, Dried Bovine Colostrum, Animal and Vegetable Fat (preserved with BHA and BHT), Dextrose, Lecithin, L-Lysine Monohydrochloride, Extracted Citric Acid Presscake, DL-Methionine, Vitamin A Acetate, Vitamin D₃ Supplement, Vitamin E Supplement, Ascorbic Acid, Choline Chloride, Vitamin B₁₂ Supplement, Menadione Sodium Bisulfite Complex (Source of Vitamin K Activity), Niacin Supplement, Calcium Pantothenate, Pyridoxine Hydrochloride, Riboflavin Supplement, Folic Acid, Thiamine Mononitrate, Biotin, Ethylenediamine Dihydriodide, Calcium Carbonate, Dicalcium Phosphate, Magnesium Sulfate, Cobalt Sulfate, Ferrous Sulfate, Selenium Yeast, Brewer's Dried Yeast, Manganese Sulfate, Zinc Sulfate, Manganese Methionine Complex, Iron Amino Acid Complex, Zinc Methionine Complex, Mono and Diglycerides of Edible Fats or Oils, Natural and Artificial Flavors.

MANAGEMENT TIPS:

1. Provide all lambs and goat kids with colostrum milk during the first 24 hours to provide concentrated nutrients and passive immunity to disease. Provide 10% of animal's bodyweight split into 3 or 4 feedings.
2. Mix and feed according to package instructions and the chart above.
3. Feed regularly and in sufficient quantity.
4. Wash bottle thoroughly after each use.
5. Provide fresh water at all times.
6. Keep lambs and goat kids in a well-ventilated, draft free area to avoid sudden temperature changes with supplemental heat if needed.
7. A good quality grain mix and water should be offered free-choice after lambs or goat kids reach about one week of age.
8. Good quality roughage should be offered free-choice when lambs or goat kids reach about 3 weeks of age.
9. Lambs can be weaned when doubled their birthweight or are consuming 0.5 pounds of dry feed per head per day. Goat kids can be weaned at 2.5 times their birthweight.
10. It is recommended that orphan lambs and goat kids be vaccinated weekly for 3-4 weeks with a *Clostridium perfringens* type CD&T Toxoid.

Manufactured for



Premier 1 Supplies
2031 300th St, Washington, IA 52353
800-282-6631
premier1supplies.com

MADE IN THE U.S.A.
Net Weight 25 lb (11.34 kg)

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed.

For further information on feeding milk replacer and colostrum needs contact premier1supplies.com.