

**Clostridium Perfringens
Types C & D-
Tetanus Toxoid**

Bar Vac® CD/T

250 mL/50 Cattle Doses or
125 Sheep/Goat Doses

Indications: Recommended for the vaccination of healthy, susceptible sheep, goats and cattle against enterotoxemia and tetanus caused by the toxins of *Clostridium perfringens* Types C and D and *Clostridium tetani*. Although *Cl. perfringens* Type B is not a significant problem in the U.S.A., immunity may be provided against the beta and epsilon toxins elaborated by *Cl. perfringens* Type B. This immunity is derived from the combination of Type C (beta) and Type D (epsilon) fractions.

Composition: Prepared from cultures of the organisms listed. Alum precipitated.

Dosage:

Cattle: Using aseptic technique, inject 5 mL subcutaneously. Repeat in 21 to 28 days and once annually.

Sheep and Goats: Using aseptic technique, inject 2 mL subcutaneously. Repeat in 21 to 28 days and once annually.

Precautions: Store out of direct sunlight at 35-45°F (2-7°C). Avoid freezing. Shake well before using. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur.

Antidote: Administer epinephrine.

Boehringer Ingelheim Vetmedica, Inc.
St. Joseph, Missouri 64506 U.S.A.
U.S. Veterinary License No. 124

Code 120-321; 20302-04

