

Vaccines for sheep and goats

Susan Schoenian, Sheep & Goat Specialist, University of Maryland Extension [last updated 4.19.2012]

| Vaccine | Manufacturer | Approved Species | Specs | Sheep/Goat Dosage | Slaughter Withdrawal |
|-------------------------------------|--------------------------|---|---|---|----------------------|
| Bar-Vac® CD-T | Boehringer | Cattle Sheep Goats | Vaccine for immunizing sheep, goats, and cattle against tetanus and overeating disease | 2 ml SQ Repeat in 21 to 28 days Repeat annually | 21 days |
| C & D Antitoxin | Boehringer- Ingleheim | Cattle Calves Sheep Suckling lambs Goats Baby pigs | Vaccine for prevention (lasting 3 weeks) and treatment of enterotoxemia | Sheep: 10 ml IM or SQ Suckling lambs: 3 ml For treatment, double above doses | 21 days |
| C & D Toxoid | Colorado Serum | Cattle Sheep | For vaccination against enterotoxemia caused by the toxins of Clostridium perfringens type C and D. | 2 ml SQ Repeat in 21 to 28 days Repeat annually | 21 days |
| Case-Bac™ | Colorado Serum | Sheep Goats | Vaccine to aid in the prevention and control of Caseous lymphadenitis | 2 ml SQ (auxillary space) Repeat full 2 ml dose in 4 weeks. Annual booster | 21 days |
| Campylobacter fetus bacterin | Colorado Serum | Sheep | Protect ewes against vibrio abortion | 5 ml SQ on side of neck or rib cage shortly before exposure to ram, 2 nd dose 60 to 90 days later, repeat annually | 21 days |
| Caseous D-T | Colorado Serum | Sheep | Vaccine against Caseous Lymphadenitis, overeating disease, and lock jaw | 2 ml SQ to sheep of any age, repeat in 4 weeks, annual booster recommended | 21 days |

| Vaccine | Manufacturer | Approved Species | Specs | Sheep/Goat Dosage | Slaughter Withdrawal |
|---|-----------------|---|---|--|----------------------|
| Chlamydia Psittaci Bacterin | Colorado Serum | Sheep Goats | Protect ewes against ovine enzootic abortion | 2 ml SQ in the top of the neck 60 days prior to breeding, repeat 2 ml dose in 30 days, revaccinate annually just prior to breeding | 60 days |
| Clostratox BCD | Novartis | Cattle Sheep | Aid in the prevention of Enterotoxemia caused by Clostridium perfringens Types C and D. The combination of these fractions may protect against Type B | Feeder lambs: 10 ml SQ Double dosage for treatment | 21 days |
| Clostridium perfringens type C & D antitoxin | Colorado Serum | Suckling calves Feeder calves Suckling pigs Suckling lambs Suckling goats | Multivalent antitoxin to treat and prevent enterotoxemia caused by types B, C, and D | Suckling lambs and kids: 5 ml Suckling feeder lambs: 10 ml | 21 days |
| Covexin®-8 | Schering-Plough | Cattle Sheep | For the vaccination of healthy cattle and sheep against diseases caused by <i>Clostridium chauvoei</i> , <i>Cl. septicum</i> , <i>Cl. novyi</i> Type B, <i>Cl. haemolyticum</i> (known elsewhere as <i>Cl. novyi</i> Type D), <i>Cl. tetani</i> and <i>Cl. perfringens</i> Types C and D. | 5 ml SQ, followed by 2 ml dose in six weeks. Repeat annually with 2 ml dose. | 21 days |
| Ovine Ecthyma vaccine Live virus. | Colorado Serum | Sheep Goats | Protect healthy sheep and goats against soremouth | Apply to wool-free, scarified area Do not use within 24 hours of dipping or spraying | 21 days |
| Essential 3 + T | Colorado Serum | Cattle Swine Sheep Goats | Aid in prevention of enterotoxemia caused by Cl. Perfringens Type B, C, and D and tetanus | 2 ml SQ Repeat in 3-4 weeks Repeat annually | 21 days |

| Vaccine | Manufacturer | Approved Species | Specs | Sheep/Goat Dosage | Slaughter Withdrawal |
|--|----------------------|---|--|--|----------------------|
| Pasteurella (pneumonia) | Colorado Serum | Cattle Sheep Goats | Aid in prevention of pneumonia caused by Mannheimia haemolytica and Pasteurella multocida | 2 ml SQ Repeat in 2-4 weeks | 21 days |
| Rabies vaccine Imrab 3 Imrab Large Animal | Merial | Cattle Sheep Horses Dogs Cats | Killed virus recommended for vaccination for prevention of disease due to rabies. | 2 ml SQ Repeat in 1 year and every 3 years thereafter | 21 days |
| Tetanus Antitoxin | Colorado Serum | Horses Cattle Sheep Swine Goats | For use whenever a non-immunized animal, or one whose immune status is unknown, suffers a deep penetrating wound that has or may become contaminated with soil. It provides quick but short-term protection. | Administer 1500 units (5 ml) SQ or IM for 7-14 days of passive immunity For treatment, administer 3000-15000 units. Administer 2 nd dose in 7 days | 21 days |
| Tetanus Toxoid | Colorado Serum | Sheep Goats Swine Cattle Horses | Vaccine for establishing immunity against tetanus | 2 doses of 1 ml per 100 lbs. IM each 30 days apart | 21 days |
| Ultrabac 8 | Pfizer Animal Health | Cattle Sheep | Aid in preventing disease caused by Cl. chauvoei, septicum, haemolyticum, novyi, sordellii and perfringens types C & D. Immunity may be provided against type B. | 2.5 ml SQ Repeat in 4-6 weeks Repeat annually | 21 days |
| Ultrabac CD | Pfizer Animal Health | Cattle Sheep | For the immunization of healthy cattle and sheep against Enterotoxemia Enteritis caused by Cl. perfringens types C & D. | 1 ml IM or SQ Repeat in 4-6 weeks Repeat annually | 21 days |

| Vaccine | Manufacturer | Approved Species | Specs | Sheep/Goat Dosage | Slaughter Withdrawal |
|--------------------------------|----------------------|--------------------------|---|---|----------------------|
| UltraChoice 8 | Pfizer Animal Health | Cattle Sheep | Aid in preventing diseases caused by Cl. chauvoei, septicum, novyi, sordellii, haemolyticum and perfringens types C and D | 1 ml SQ in neck Repeat in 4-6 weeks Repeat annually | 21 days |
| Vision® CD-T with Spur® | Intervet | Cattle Sheep Goats | Aid in the prevention of enterotoxemia caused by Clostridium perfringens types C & D | 2 ml SQ Repeat dose in 3 to 4 weeks | 21 days |

Source: product labels