

154025  
09M0420

**INDICATIONS:** An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves. For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

**DOSAGE AND ADMINISTRATION:**  
**5-Day Treatment Protocol**

Daily Dosage: 10 mg amprolium/kg  
(10 mg per 2.2 lb body weight)

**21-Day Prevention Protocol**

Daily Dosage: 5 mg amprolium/kg  
(5 mg per 2.2 lb body weight)

**USE DIRECTIONS:** There are **96 mg** of amprolium in every **1 mL** of CORID 9.6% Solution.

1 fl oz = 29.57 mL      8 fl oz = 1/2 pint (236.56 mL)  
1 fl oz = 2 measuring tablespoons      16 fl oz = 1 pint (473.12 mL)

**IN DRINKING WATER**

Mix CORID daily with fresh drinking water.

**5-DAY TREATMENT:** Add CORID 9.6% Oral Solution to drinking water at the rate of 16 fl oz/100 gal. At the usual rate of water consumption this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

**21-DAY PREVENTION:** During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add CORID 9.6% Oral Solution to drinking water at the rate of 8 fl oz/100 gal. At usual rates of water consumption this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 21 days.

© CORID and the CATTLE HEAD logo are registered trademarks of Merial Limited.  
1022-2220-02  
Rev. 02-2008

Lot No & Exp Date ▼

**AS A DRENCH**

CORID drench solutions may be stored in a clean, closed, labeled container for up to 3 days.

**5-DAY TREATMENT:** Add 3 fl oz CORID 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2 lb) body weight. Give daily for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

**21-DAY PREVENTION:** During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1 1/2 fl oz of CORID 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21 days.

**RESIDUE WARNINGS:** Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**WARNING: NOT FOR HUMAN USE. Keep this and all drugs out of the reach of children.** The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. To report adverse reactions in users, to obtain more information, or to obtain a MSDS, contact Merial at 1-888-637-4251.

**PRECAUTIONS: FOR ORAL USE IN ANIMALS ONLY. MAY CAUSE EYE IRRITATION.** For irritation, flush with plenty of water; get medical attention.  
Restricted Drug (California) - Use only as directed.

**STORAGE:** Store between 5° - 25° C (41° - 77° F) with brief excursions to 40° C. Benzoic acid 0.1% added as preservative

Technical Assistance: 1-888-637-4251  
www.corid.com

