

Three 1 gram pouches for a 3-day treatment.

Each pouch treats 10 lbs.

Intervet

# safe-guard®

(fenbendazole)



# Canine Dewormer

For dogs only, 6 weeks and older.

**Drug Facts:**  
Active ingredients (in each dosage unit): Fenbendazole Granules 22.2% (222 mg/g)

**Uses: For the treatment and control of:**

- **Tapeworms**  
(*Taenia pisiformis*)
- **Roundworms**  
(*Toxocara canis*, *Toxascaris leonina*)
- **Hookworms**  
(*Ancylostoma caninum*, *Uncinaria stenocephala*)
- **Whipworms**  
(*Trichuris vulpis*)



Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

**HUMAN WARNING:**  
KEEP OUT OF REACH OF CHILDREN.

446100-C  
IC-30205/G

Net Weight: 3 grams

Intervet

## safe-guard®

(fenbendazole)

**Directions:** The daily dose for Safe-Guard® Canine is 50 mg/kg (22.7 mg/lb) of body weight. You should weigh your dog to make sure you are using the right dose; i.e. correct gram amount and appropriate number of pouches.

**Your dog must be treated with the following dose for 3 days in a row.**

Dog Weight (lbs)	Pouch Size (Daily Dose)
10	1 gram
11 to 20	2 gram
21 to 30*	1 gram + 2 gram
31 to 40	4 gram
41 to 50*	1 gram + 4 gram
51 to 60*	2 gram + 4 gram
61 to 80*	Two 4 gram
Over 80*	* Combine pouches to obtain recommended daily dosage.

Mix daily dose with a small amount of the usual food. Make sure your dog eats all the medicated food. You may need to moisten dry dog food to aid mixing.

Repeat the dose for 3 days in a row.



**Recommended Deworming Schedule:**  
Schedule may vary depending on the climate where you live and the activity of your dog. The following schedule should be used as a general guideline.

- Weaned pups (6-8 weeks of age) should be dewormed at 6, 8, 10 and 12 weeks of age.
- Treat the dam at the same time as the puppies.
- Dogs over 6 months of age should be dewormed at least twice each year.

**Diagnosis of Parasites:** Consult your veterinarian for help in the diagnosis, treatment and control of parasitism. Specific diagnoses require laboratory testing. Dogs can become infected with several different kinds of tapeworms. Safe-Guard® Canine only kills *Taenia* species of tapeworms. If you continue to see tapeworm segments in your dog's stool after treatment with Safe-Guard® Canine, consult your veterinarian.

Roundworms and hookworms may pose serious human health risks; regular deworming may help reduce those risks.

**Product Use:** Safe-Guard® Canine is safe for use in puppies 6 weeks or older and adult dogs, including pregnant bitches. Do not deworm a dog or puppy that is sick. Consult your veterinarian for diagnosis of the illness.

**Adverse Reactions:** In U.S. clinical studies 3 of 240 dogs (about 1% of the treated dogs) had vomiting associated with the use of the product.

**Storage:** Store at controlled room temperature (59-86°F).

**Questions? Comments?**

To obtain product information including material safety data sheets (MSDS) or to report a suspected adverse reaction, call 1-800-441-8272.

Distributed by:  
INTERVET INC., Millsboro, DE

NADA #121-473, Approved by FDA

Made in Canada

[www.Safe-Guard-For-Dogs.com](http://www.Safe-Guard-For-Dogs.com)



Three 2 gram pouches  
for a 3-day treatment.

Each pouch treats 20 lbs.

Intervet  
**safe-guard**<sup>®</sup>  
(fenbendazole)



# Canine Dewormer

For dogs only, 6 weeks and older.

**Drug Facts:**  
Active ingredients (in each dosage unit): Fenbendazole Granules 22.2% (222 mg/g)

**Uses: For the treatment and control of:**

- **Tapeworms**  
(*Taenia pisiformis*)
- **Roundworms**  
(*Toxocara canis*, *Toxascaris leonina*)
- **Hookworms**  
(*Ancylostoma caninum*,  
*Uncinaria stenocephala*)
- **Whipworms**  
(*Trichuris vulpis*)



Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

**HUMAN WARNING:**  
KEEP OUT OF REACH OF CHILDREN.

Intervet  
**safe-guard**<sup>®</sup>  
(fenbendazole)


**Directions:** The daily dose for Safe-Guard<sup>®</sup> Canine is 50 mg/kg (22.7 mg/lb) of body weight. You should weigh your dog to make sure you are using the right dose; i.e. correct gram amount and appropriate number of pouches.

**Your dog must be treated with the following dose for 3 days in a row.**

Dog Weight (lbs)	Pouch Size (Daily Dose)
10	1 gram
11 to 20	2 gram
21 to 30*	1 gram + 2 gram
31 to 40	4 gram
41 to 50*	1 gram + 4 gram
51 to 60*	2 gram + 4 gram
61 to 80*	Two 4 gram
Over 80*	* Combine pouches to obtain recommended daily dosage.

Mix daily dose with a small amount of the usual food. Make sure your dog eats all the medicated food. You may need to moisten dry dog food to aid mixing.

Repeat the dose for 3 days in a row.



**Recommended Deworming Schedule:**

Schedule may vary depending on the climate where you live and the activity of your dog. The following schedule should be used as a general guideline.

- Weaned pups (6-8 weeks of age) should be dewormed at 6, 8, 10 and 12 weeks of age.
- Treat the dam at the same time as the puppies.
- Dogs over 6 months of age should be dewormed at least twice each year.

**Diagnosis of Parasites:** Consult your veterinarian for help in the diagnosis, treatment and control of parasitism. Specific diagnoses require laboratory testing. Dogs can become infected with several different kinds of tapeworms. Safe-Guard<sup>®</sup> Canine only kills *Taenia* species of tapeworms. If you continue to see tapeworm segments in your dog's stool after treatment with Safe-Guard<sup>®</sup> Canine, consult your veterinarian.

Roundworms and hookworms may pose serious human health risks; regular deworming may help reduce those risks.

**Product Use:** Safe-Guard<sup>®</sup> Canine is safe for use in puppies 6 weeks or older and adult dogs, including pregnant bitches. Do not deworm a dog or puppy that is sick. Consult your veterinarian for diagnosis of the illness.

**Adverse Reactions:** In U.S. clinical studies 3 of 240 dogs (about 1% of the treated dogs) had vomiting associated with the use of the product.

**Storage:** Store at controlled room temperature (59-86°F).

**Questions? Comments?**

To obtain product information including material safety data sheets (MSDS) or to report a suspected adverse reaction, call 1-800-441-8272.



Distributed by:  
INTERVET INC., Millsboro, DE  
NADA #121-473, Approved by FDA  
Made in Canada

[www.Safe-Guard-For-Dogs.com](http://www.Safe-Guard-For-Dogs.com)



Three 4 gram pouches for a 3-day treatment.

Each pouch treats 40 lbs.

# safe-guard® (fenbendazole)



# Canine Dewormer

For dogs only, 6 weeks and older.

Drug Facts:  
Active ingredients (in each dosage unit): Fenbendazole Granules 22.2% (222 mg/g)

### Uses: For the treatment and control of:

- **Tapeworms**  
(*Taenia pisiformis*)
- **Roundworms**  
(*Toxocara canis*, *Toxascaris leonina*)
- **Hookworms**  
(*Ancylostoma caninum*, *Uncinaria stenocephala*)
- **Whipworms**  
(*Trichuris vulpis*)



Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

**HUMAN WARNING:**  
KEEP OUT OF REACH OF CHILDREN.

446400-C  
IC-30218/G

Net Weight: 12 grams

## safe-guard® (fenbendazole)


**Directions:** The daily dose for Safe-Guard® Canine is 50 mg/kg (22.7 mg/lb) of body weight. You should weigh your dog to make sure you are using the right dose; i.e. correct gram amount and appropriate number of pouches.

**Your dog must be treated with the following dose for 3 days in a row.**

Dog Weight (lbs)	Pouch Size (Daily Dose)
10	1 gram
11 to 20	2 gram
21 to 30*	1 gram + 2 gram
31 to 40	4 gram
41 to 50*	1 gram + 4 gram
51 to 60*	2 gram + 4 gram
61 to 80*	Two 4 gram
Over 80*	* Combine pouches to obtain recommended daily dosage.

Mix daily dose with a small amount of the usual food. Make sure your dog eats all the medicated food. You may need to moisten dry dog food to aid mixing.

Repeat the dose for 3 days in a row.



**Recommended Deworming Schedule:**  
Schedule may vary depending on the climate where you live and the activity of your dog. The following schedule should be used as a general guideline.

- Weaned pups (6-8 weeks of age) should be dewormed at 6, 8, 10 and 12 weeks of age.
- Treat the dam at the same time as the puppies.
- Dogs over 6 months of age should be dewormed at least twice each year.

**Diagnosis of Parasites:** Consult your veterinarian for help in the diagnosis, treatment and control of parasitism. Specific diagnoses require laboratory testing. Dogs can become infected with several different kinds of tapeworms. Safe-Guard® Canine only kills *Taenia* species of tapeworms. If you continue to see tapeworm segments in your dog's stool after treatment with Safe-Guard® Canine, consult your veterinarian.

Roundworms and hookworms may pose serious human health risks; regular deworming may help reduce those risks.

**Product Use:** Safe-Guard® Canine is safe for use in puppies 6 weeks or older and adult dogs, including pregnant bitches. Do not deworm a dog or puppy that is sick. Consult your veterinarian for diagnosis of the illness.

**Adverse Reactions:** In U.S. clinical studies 3 of 240 dogs (about 1% of the treated dogs) had vomiting associated with the use of the product.

**Storage:** Store at controlled room temperature (59-86°F).

### Questions? Comments?

To obtain product information including material safety data sheets (MSDS) or to report a suspected adverse reaction, call **1-800-441-8272**.



Distributed by:  
INTERVET INC., Millsboro, DE

NADA #121-473, Approved by FDA

Made in Canada

[www.Safe-Guard-For-Dogs.com](http://www.Safe-Guard-For-Dogs.com)

