



Liquamycin[®]
LA-200[®]
(oxytetracycline injection)

Antibiotic

*Each mL contains 200 mg
of oxytetracycline base as
oxytetracycline dihydrate.*

*For the treatment of disease
in beef cattle; dairy cattle; calves,
including preruminating (veal)
calves; and swine*

For animal use only

Liquamycin[®]

LA-200[®]

(oxytetracycline injection)

Cattle Dosage Guide

At the first signs of pneumonia or pinkeye,* administer a single dose of Liquamycin LA-200 subcutaneously according to the following weight categories.**

Animal Weight (lb)	Number of mL or cc	Animal Weight (lb)	Number of mL or cc
100	4.5	700	31.5
200	9.0	800	36.0
300	13.5	900	40.5
400	18.0	1000	45.0
500	22.5	1100	49.5
600	27.0	1200	54.0

* See package insert for dosing instructions for other indicated diseases and full product information.

** Do not administer more than 10 mL at any one injection site (1–2 mL per site in small calves).

Discontinue treatment at least 28 days prior to slaughter.

Swine Dosage Guide

At the first signs of pneumonia,* administer Liquamycin LA-200 by deep intramuscular injection according to the following weight categories.**

Animal Weight (lb)	Number of mL or cc	Animal Weight (lb)	Number of mL or cc
10	0.5	175	7.9
25	1.1	200	9.0
50	2.3	225	10.1
75	3.4	250	11.3
100	4.5	275	12.4
125	5.6	300	13.5
150	6.8	325	14.6

* See package insert for dosing instructions for other indicated diseases and full product information.

** Do not administer more than 5 mL at any one injection site.

Discontinue treatment at least 28 days prior to slaughter.

7960000

10-4983-00-2

Made in Brazil

U.S. Patent No. 4,018,889

Liquamycin LA-200 (oxytetracycline injection) is a sterile, preconstituted solution of the broad-spectrum antibiotic oxytetracycline.

Caution: When administered to cattle, muscle discoloration may necessitate trimming of the injection site(s) and surrounding tissues during the dressing procedure.

Warnings: Discontinue treatment at least 28 days prior to slaughter of cattle and swine. Milk taken from animals during treatment and for 96 hours after the last treatment must not be used for food. Rapid intravenous administration may result in animal collapse. Oxytetracycline should be administered intravenously over a period of at least 5 minutes.

Precautions: Exceeding the highest recommended level of drug per lb of body weight per day, administering more than the recommended number of treatments, and/or exceeding 10 mL subcutaneously per injection site in adult beef and dairy cattle, and 5 mL intramuscularly per injection site in adult swine, may result in antibiotic residues beyond the withdrawal period.

Refer to Package Insert for Complete Directions

Storage: Store at room temperature 15°–30°C (59°–86°F). Keep from freezing.

Restricted Drug (California)—Use Only as Directed

Not for Human Use

Laboratórios Pfizer Ltda. - Animal Health Division
Av. Monteiro Lobato, 2270, Guarulhos, São Paulo,
Brasil - CNPJ nº 46.070.868/0001-69

Licenciado no Ministério da Agricultura sob o
nº 801/78, em 29/11/78

Licensed in the Ministry of Agriculture under
nº 801/78, on 11/29/78



Distributed by:



Pfizer Animal Health

Div. of Pfizer Inc
NY, NY 10017