

Intervet

safe-guard[®]
(fenbendazole)



Dewormer

for Beef & Dairy Cattle and Goats

Suspension 10% (100 mg/mL)

RESIDUE WARNINGS: Cattle must not be slaughtered within 8 days following treatment. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Goats must not be slaughtered for food within 6 days following treatment. Because a withdrawal time in milk has not been established, do not use in lactating goats. For dairy cattle, there is no milk withdrawal period.

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

Keep this and all medication out of the reach of children.

1,000 mL (33.8 fl oz)

697510-E

INDICATIONS:

Beef and Dairy Cattle - 2.3 mg/lb (5 mg/kg) body weight for the removal and control of: Lungworm (*Dichyoaulus viviparus*); Stomach worm (adults): brown stomach worm (*Ostertagia ostertagi*); Stomach worms (adults and fourth stage larvae): barberpole worm (*Haemonchus contortus* and *H. placei*), and small stomach worm (*Trichostrongylus axei*); Intestinal worms (adults and fourth stage larvae): hookworm (*Bunostomum phlebotomum*), threadnecked intestinal worm (*Nematodirus helvetianus*), small intestinal worm (*Cooperia punctata* and *C. oncophora*), bankrupt worm (*Trichostrongylus colubriformis*) and nodular worm (*Oesophagostomum radiatum*).

Goats - 2.3 mg/lb (5 mg/kg) body weight for the removal and control of: Stomach worms (adults): *Haemonchus contortus* and *Teladorsagia circumcincta*.

DIRECTIONS:

Determine the proper dose according to estimated body weight. Administer orally. The recommended dose of 2.3 mg/lb (5 mg/kg) of body weight is achieved when 2.3 mL of the drug are given for each 100 lb body weight.

EXAMPLES:

Cattle:

Body Weight	Amount
100 lb	2.3 mL
200 lb	4.6 mL
300 lb	6.9 mL
400 lb	9.2 mL
500 lb	11.5 mL
1000 lb	23.0 mL
1500 lb	34.5 mL

Goats:

Body Weight	Amount
25 lb	0.6 mL
50 lb	1.2 mL
75 lb	1.7 mL
100 lb	2.3 mL
125 lb	2.9 mL

Under conditions of continued exposure to parasites, retreatment may be needed after 4 to 6 weeks.

Manufactured by: DPT Laboratories, San Antonio, TX 78215

Distributed by: Intervet Inc., Millsboro, DE 19966

Restricted drug (California). Use only as directed. Store at or below 25°C (77°F).

Protect from freezing.

Shake well before use.

For Use in Animals Only.



LOT NUMBER:

AMCR

EXPIRATION DATE:

12/12

697515-E

NADA # 128-620, Approved by FDA

