



Ready To Use

# QuickBayt™ Fly Bait

For the control of nuisance flies in and around livestock facilities and stables.

- ✓ Fast acting
- ✓ Ready-to-use as a scatter bait, paint-on or in bait stations
- ✓ Contains a specially formulated mixture of attractants in extruded pellet form

**ACTIVE INGREDIENTS:**

Imidacloprid.....	0.50%
Z-9-Tricosene.....	0.10%

**OTHER INGREDIENTS:**..... 99.40%

**TOTAL:**.....100.00%

**STOP—Read the label before use.  
Keep Out of Reach of Children.**

## CAUTION

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed, absorbed through skin, or inhaled. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco.

#### FIRST AID

<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>• Do not give anything to an unconscious person.</li> </ul>
<b>IF ON SKIN:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

#### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For product questions, to report adverse reactions, or for a copy of the Safety Data Sheet (SDS), call Elanco Product & Veterinary Support at 1-800-428-4441.

**Note to Physician:** No specific antidote is available. Treat the patient symptomatically.

#### ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

#### PRO FACTS

- Fast acting
- Ready-to-use as scatter bait, paint-on or in bait stations
- Contains a specially formulated mixture of attractants in extruded pellet form
- Controls flies resistant to organophosphates and carbamates

**QUICKBAYT FLY BAIT**

- Ready-to-use bait containing 0.5% imidacloprid
- Contains the attractant Z-9-Tricosene
- Contains Bitrex™ bittering agent

**CONTROLS**

- Nuisance flies

**COVERAGE** 1.5 oz/250 sq ft (1.5 oz = 3 Tablespoons)

**RE-APPLY** Re-apply every 7 days as needed.

**QUESTIONS??** For questions or comments, call tollfree 1-800-428-4441.



## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **IMPORTANT:** Read the entire label and Conditions of Sale before using this product.

### FLY MANAGEMENT

#### WHERE TO APPLY

**Do not place QuickBayt Fly Bait in areas accessible to animals, children and pets.**

#### Scatter Application

For use around the **outside** of:

- Agricultural production facilities like broiler houses, turkey houses, caged layer houses, feedlots, dairy barns, and swine confinement buildings
- Canneries; dairy, meat and poultry processing plants; food processing plants; warehouses; dumpsters; recycling areas and loading docks

#### Scatter Application

For use **in** walkways of:

- Caged layer houses

#### Bait Station Application

For use **inside** of:

- Horse stables
- Broiler, turkey, and caged layer houses
- Swine confinement buildings
- Dairy barns and milking parlors

#### Paint-on Application

For use **inside** and around the **outside** of:

- Broiler, turkey, and caged layer houses
- Swine confinement buildings
- Dairy barns

1.5 oz/250 sq ft (scatter bait and bait station)

**Do not place QuickBayt Fly Bait in areas accessible to animals, children and pets.**

Apply the bait in many locations. Avoid cool or windy locations because flies prefer warm, motionless sites for resting. Time applications to begin at the start of the season before fly populations have reached their peak.

**Scatter Application:** Scatter the bait directly from the container onto dry level surfaces so that the individual granules lie near each other without forming small piles.

**Bait Station Application:** Place in any fly bait station. Use 1 bait station to cover approximately 250 sq ft, and add up to 1.5 oz of bait per station. Secure bait stations at least 4 feet above the ground.

**Paint-on Application:** Mix 1.5 oz bait with 1 fl oz warm water (3 lbs bait in 1 quart water) and stir thoroughly. Let stand for about 15 minutes until a paste consistency suitable for painting has formed. Apply the paste with a brush to surfaces where flies rest avoiding surfaces that are dusty.

#### RESTRICTIONS



- Do not place in areas accessible to food producing animals, children and pets.
- Do not allow contact with food or water.
- Use environmental controls, such as proper light management.
- Do not apply directly onto milking equipment.
- Do not apply inside food processing plants or restaurants.

#### MANAGEMENT TIPS

**For Best Results:** Use the bait as part of an overall Integrated Pest Management program.

- Use proper sanitation and manure management.
- Use environmental controls, such as proper light management.
- Consult with Cooperative Extension.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.	
<b>Pesticide Storage</b>	<ul style="list-style-type: none"> <li>• Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, or feed.</li> <li>• Store in original containers and out of the reach of children, preferably in a locked storage area.</li> <li>• Handle an open container in a manner as to prevent spillage.</li> </ul>
<b>Pesticide Disposal</b>	<ul style="list-style-type: none"> <li>• Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.</li> </ul>
<b>Container Disposal</b>	<ul style="list-style-type: none"> <li>• Completely empty container. Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Offer for reconditioning, if appropriate. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.</li> <li>• Dispose of container by offering for recycling or reconditioning, or at an approved waste disposal facility or by incineration if allowed by state and local authorities.</li> <li>• Do not use container in connection with food, feed, or drinking water.</li> </ul>
<b>Spills</b>	<ul style="list-style-type: none"> <li>• If the container is leaking or material is spilled, carefully sweep up spilled bait and wash exposed area.</li> <li>• In spill or leak incidents, keep unauthorized people away.</li> </ul>

**Conditions of Sale:**

The directions on this label are believed to be adequate and must be followed carefully. This product has been tested under different environmental conditions similar to those that are ordinary and customary for use of the product. However, it is impossible to eliminate all risks associated with the use of this product. Failure to follow label directions may cause injury to animals, humans and damage to the environment. Insufficient control of pests, injury or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Elanco. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

**Limited Warranty and Limitation of Damages:** Elanco warrants that this material conforms to the chemical description on the label. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ELANCO MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL, and no agent of Elanco is authorized to make or modify any warranties beyond those contained herein. Any damages arising from breach of this warranty shall be limited to direct damages and shall not include special, incidental or consequential damages resulting from the use of handling of this product.

EPA Reg. No. 101563-215-11556

Distributed by Elanco US Inc. 2500 Innovation Way,  
Greenfield, IN 46140

Revised: June 2023

The QuickBayt trademark is owned by Bayer and is used under license. Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.  
© 2023 Elanco or its affiliates

